

Claw™

HERBICIDE

For the control of barley grass and annual grass weeds
in PASTURE and SPORTS TURF

Active Ingredient: 500 g/litre ethofumesate in the form
of a suspension concentrate

PLEASE READ THE LABEL COMPLETELY BEFORE USE

HAZARD CLASSIFICATIONS: 9.1B, 9.2A

WARNING: ECOTOXIC

Toxic to aquatic organisms. Very toxic to some plant species.

PRECAUTIONS

APPROVED HANDLER: This product must be under the control of an approved handler during use.

STORAGE: Keep out of reach of children. Store in the original, tightly closed container, in a secure place away from foodstuffs. Do not store in direct sunlight. Storage of certain quantities of this product trigger specific storage requirements (Refer to HazNote). When stored appropriately this product should show no significant degradation for two years from the date of manufacture. Contact your supplier for further information about the use of product that is older than this.

PERSONAL PROTECTION: Avoid contact with eyes and skin. Avoid inhalation of spray mist. When mixing or applying, wear overalls, boots, gloves and eye protection. Clean protective equipment daily after work. Do not eat, drink or smoke while using. Remove protective clothing and wash hands and face thoroughly before meals and after work. **USE:** Apply with well maintained and calibrated spray equipment.

DISPOSAL: Dispose of this product only by using in accordance with this label, or at an approved landfill, or other approved facility. Triple rinse container and add rinsate to the spray tank. Dispose of container to an approved landfill or agricultural recycling scheme. Do not burn. Do not use container for any other purpose.

ENVIRONMENTAL: Use this product carefully. Do not contaminate aquatic environments with product, spray drift or empty container. Spray drift or equipment contamination may cause serious damage to desirable plants. Do not allow spray drift outside the target area to occur.

Emergency Response

In a transport emergency call **111, Police or Fire Brigade**. For specialist advice in an emergency only, call **0800 CHEMCALL (0800 243 6225)** (24 hrs).

Spillage

Exclude non-essential people from the area. Wear protective clothing and prevent material from entering waterways. Absorb spill with inert material such as soil, sand, etc. Place in waste container. Wash area with water, and absorb with further inert material. Dispose of safely to approved landfill.

Safety Data Sheet

Additional information is provided in the safety data sheet. Freephone 0800 ORIONCP (0800 674 6627) or visit www.orioncp.co.nz.

GENERAL INFORMATION

Claw is a selective herbicide for the control of barley grass and annual grass weeds in pasture and sports turf. Best results are obtained when weeds are not under stress.

Weeds Controlled. Susceptible: Redroot, fumitory, cleavers, chickweed, spurrey, wild portulaca, summer grass, annual poa, sow thistle. **Moderately Susceptible:** Fathen, black nightshade, wireweed, willow weed, speedwells, banyard grass, wild oats, combind, annual nettle, shepherds purse. **Resistant:** Wild radish, dead nettle, galinsoga, ryegrasses, groundsel, perennial weeds. **Controlled Pre-Emergence Only:** Cornbind, annual nettle, shepherds purse.

APPLICATION METHODS

Non-reinforced PVC tubing may soften and swell after prolonged contact. Use conventional ground spraying equipment. Clean sprayer thoroughly with water and detergent.

COMPATIBILITY

Do NOT mix with any wetting agents or surfactants. May be tank mixed with products containing phenmedipham and desmedipham. Mixing with phenoxy herbicides for use on pasture is NOT recommended.

RAINFALL

Do not apply if rain is expected within 3 hours of spraying.

WITHHOLDING PERIODS

Do NOT graze stock on treated areas for 28 days.

WARRANTY

This formulation is warranted to contain the specified active ingredient within accepted

DIRECTIONS FOR USE

Crop	Weeds Controlled	Rate per Hectare	Rate for Gun Application	Critical Comments
Pasture	Barley grass	4 litres/ha in 200 – 400 litres water per ha	400 ml/100 litres water	Apply after main germination, usually from April to July. Evenly grazed pasture 3 - 5cm high will help achieve optimum results. Raw organic matter will reduce effectiveness on stock camps. Temporary suppression of white clover will occur.
Redbeet – beetroot		Pre-sowing and Pre-emergence: 3 - 4 litres/ha in 200 – 300 litres water per ha Post-emergence: 4 litres/ha in 200 - 300 litres water per ha		Apply before weed emergence and before or as soon as possible after drilling. The seedbed needs to be fine, firm and moist. Use the higher rate over dry and cloddy soil and incorporate to 3 - 5 cm. Apply to weeds at the cotyledon to 2 leaf stage.
Turf – ryegrass (do not apply to fine turf species)	Annual poa Summer, banyard and bristle grasses	4 litres/ha in 200 – 400 litres water per ha	400 ml/100 litres water	Treat the same as for barley grass. Delay mowing for 4 - 5 days after spraying. Apply just before weed emergence in September – October. After 6 - 8 weeks a repeat treatment maybe necessary. Oversow with ryegrass immediately if turf contains less than 5% ryegrass.

analytical tolerance when packed. The user bears the risk of damage resulting from factors beyond the manufacturer's control. The manufacturer declines all liability for damage resulting from improper storage or use of the product.

Registered pursuant to the ACVM Act 1997, No. P5693
See www.nzfsa.govt.nz/acvm/ for registration conditions.
Approved pursuant to the HSNO Act 1996, No. HSR000826
See www.ermanz.govt.nz for approval conditions.

™CLAW is a Trade Mark of Orion Crop Protection Ltd

Registrant: Jolyn Holdings (1991) Ltd, PO Box 568, Ashburton.
Distributor: Orion Crop Protection Ltd, Triton Technical Park, Building B, 14 – 22 Triton Drive, North Harbour, Auckland.

Batch No.

Date of manufacture



Net Contents: 5 Litres