

HARMFUL ▪ ECOTOXIC

KEEP OUT OF REACH OF CHILDREN

Gesapon[®] **20G**

INSECTICIDE

**A granular insecticide for the control
of grass grub and porina caterpillar in PASTURE
and carrot rust fly in CARROTS.**

Active Ingredient: 200 g/Kg of diazinon in the
form of a granule

PLEASE READ THE LABEL COMPLETELY BEFORE USE

HAZARD CLASSIFICATIONS: 6.1D, 6.8B, 6.9A, 9.1A, 9.2D, 9.3A, 9.4A

WARNING: TOXIC, ECOTOXIC

May be harmful if swallowed, inhaled or absorbed through the skin. May cause reproductive/development damage from repeated exposure. May cause neurological symptoms (See 'Symptoms of Poisoning' below) and nephrotoxic effects following repeated exposure at high doses. Very toxic to aquatic organisms. Harmful to soil organisms. Very toxic to terrestrial vertebrates and to terrestrial invertebrates.

PRECAUTIONS

APPROVED HANDLER: This product must be under the control of an approved handler during use.
STORAGE: Keep out of reach of children. Store in the original packaging, in a cool, dry, dark, well ventilated and secure place away from feed or foodstuffs. Storage of certain quantities of this product trigger specific storage requirements (Refer to HazNote). When stored appropriately this product should show no significant degradation for two years from the date of manufacture. Contact your supplier for further information about the use of product that is older than this.

PERSONAL PROTECTION: Avoid contact with eyes and skin. Avoid inhalation of dust. When handling, mixing or applying, wear overalls, boots, gloves and eye protection. Clean protective equipment daily after work. Do not eat, drink or smoke while using. Remove protective clothing and wash hands and face thoroughly before meals and after work.

USE: Apply with well maintained and calibrated equipment. Diazinon may be decomposed by acidic fertilisers. Use on the day of mixing.

DISPOSAL: Dispose of this product only by using in accordance with this label, or at an approved landfill, or other approved facility. Do not burn. Dispose of packaging to an approved landfill. Do not burn. Do not use packaging for any other purpose.

ENVIRONMENTAL: Use this product carefully. Do not contaminate aquatic environments with product or packaging.

Symptoms of Poisoning

These may include excessive sweating, headache, weakness, faintness, giddiness, nausea, stomach pain, vomiting, small pupils, blurred vision and muscle twitching.

First Aid

If swallowed do not induce vomiting. If skin or hair contact occurs, remove contaminated clothing and flush with running water. If inhaled, immediately move to fresh air and rest. Maintain half upright position if breathing is difficult. For advice, call the National Poisons Centre **0800 POISON (0800 764 766)** (24 hrs) or a doctor immediately.

Emergency Response

In a transport emergency call **111, Police or Fire Brigade**. For specialist advice in an emergency only, call **0800 CHEMCALL (0800 243 6225)** (24 hrs).

Spillage

Exclude non-essential people from the area. Wear protective clothing. Moistened to prevent dusting and sweep up. Place in a waste container. Dispose of safely to an approved landfill.

Safety Data Sheet

Additional information is provided in the safety data sheet. Freephone 0800 ORIONCP (0800 674 6627) or visit www.orioncp.co.nz.

GENERAL INFORMATION

Gesapon 20G is granular insecticide for the control of grass grub and porina caterpillar in pasture, and for the control of carrot rust fly in carrots.

Ingredients: Gesapon 20G contains 200g/Kg of diazinon as the active ingredient, and up to 7g/Kg of 1,2-ethanediol as a solvent.

UN 3077
 ENVIRONMENTALLY HAZARDOUS SUBSTANCE, SOLID, N.O.S. (DIAZINON 20%)
 PACKING GROUP III
 HAZCHEM CODE 2X
 ADDITIONAL INFORMATION: ORGANOPHOSPHORUS INSECTICIDE
 IN A TRANSPORT EMERGENCY, DIAL 111 POLICE OR FIRE BRIGADE.
 FOR SPECIALIST ADVICE IN AN EMERGENCY CALL 0800 CHEMCALL (0800 243 6225)



Batch No.

Date of manufacture

**DIRECTIONS FOR USE
 RATES AND TIMING OF APPLICATION**

Crop	Pest	Rate	Critical Comments
Pasture (Established)	Grass grub	11 Kg/ha	Apply as a broadcast application. For best results: Apply evenly to short pasture when grass grubs are feeding close to the surface It is most important that at least 13mm of rain should fall within one week after application Soil already moist at application may help control by encouraging shallower feeding of grubs and easier penetration of rain. Note: Penetration and activity of diazinon is reduced in heavy clay soils and in soils of high organic matter content. Control of grass grub may be less effective under these circumstances.
		5.5 Kg/ha	Apply drilled with the seed at the time of sowing.
Pasture (At sowing time)	Grass grub	11 Kg/ha	Broadcast evenly over cultivated ground. Work it into the soil to about 5cm depth during the final cultivation before sowing.
		5.5 Kg/ha	Apply as a broadcast application. For best results apply as evenly as possible and early in the autumn before excessive pasture damage has occurred. Note: In any weather Gesapon 20G will persist long enough to contact caterpillars which have stayed underground for several nights. Control of porina is not impaired by long grass or rain falling shortly after application.
Pasture	Porina caterpillar	5.5 Kg/ha	Apply as a broadcast application. For best results apply as evenly as possible and early in the autumn before excessive pasture damage has occurred. Note: In any weather Gesapon 20G will persist long enough to contact caterpillars which have stayed underground for several nights. Control of porina is not impaired by long grass or rain falling shortly after application.
Carrots	Carrot rust fly	11 Kg/ha	Apply beneath the seed at sowing. Surface band application along the row may also be needed in the late spring after 10 weeks growth, to protect until harvest.

APPLICATION METHODS

Gesapon 20G may be mixed and spread with superphosphate or lime but a more even spread is likely to be achieved by the use of granular spreading equipment with a fine application kit accessory for lower application rates. Spinner type fertiliser spreaders are not recommended. For aircraft application accurately calibrated attachments should be used to give a wider and more even swath.

WITHHOLDING PERIODS

Pasture Nil
 Note: Keep ducks, geese and hens off treated area for 7 days after application.
 Carrots 14 days


WARRANTY

This formulation is warranted to contain the specified active ingredient within accepted analytical tolerance when packed. The user bears the risk for damage resulting from factors beyond the manufacturer's control. The manufacturer declines all liability for damage resulting from improper storage or use of the product.

Registered pursuant to the ACVM Act 1997, No. P2130
 See www.nzfsa.govt.nz/acvm/ for registration conditions.
 Approved pursuant to the HSN0 Act 1996, No. HSR000175
 See www.ermanz.govt.nz for approval conditions.

®GESAPON is a Registered Trade Mark of a Syngenta Group Company.

Registrant: Orion Crop Protection Ltd,
 Triton Technical Park, Building B,
 14 – 22 Triton Drive, North Harbour, Auckland.



Net Contents: 22kg